

Measurement of the Surface Gravity of η Boo

Gerard T. van Belle¹, David R. Ciardi, Andrew F. Boden

Michelson Science Center, California Institute of Technology, Pasadena, CA 91125
gerard, bode, ciardi@ipac.caltech.edu

ABSTRACT

Direct angular size measurements of the G0IV subgiant η Boo from the Palomar Testbed Interferometer are presented, with limb-darkened angular size of $\theta_{LD} = 2.1894^{+0.0055}_{-0.0140}$ mas, which indicate a linear radius of $R = 2.672 \pm 0.028 R_{\odot}$. A bolometric flux estimate of $F_{BOL} = 22.1 \pm 0.28 \times 10^{-7}$ erg cm⁻²s⁻¹ is computed, which indicates an effective temperature of $T_{EFF} = 6100 \pm 28$ K and luminosity of $L = 8.89 \pm 0.16 L_{\odot}$ for this object. Similar data are established for a check star, HD 121860. The η Boo results are compared to, and confirm, similar parameters established by the *MOST* asteroseismology satellite. In conjunction with the mass estimate from the *MOST* investigation, a surface gravity of $\log g = 3.817 \pm 0.016$ [cm s⁻²] is established for η Boo.

Subject headings: infrared: stars, stars: fundamental parameters, techniques: interferometric, stars: individual: η Boo

1. Introduction

The bright G0 IV star η Boötis (8 Boo, HR 5235, HD 121370) is an interesting target to study given its place on the H-R diagram, and its implications upon stellar modeling. Solar-like oscillations were detected for η Boo by Kjeldsen et al. (1995, 2003). η Boo was recently observed with the Microvariability and Oscillations of STars (*MOST*) satellite (Guenther et al. 2005), a 15-cm aperture satellite observatory orbited in 2003 June by the Canadian Space Agency (Walker et al. 2003). Given that the convective envelope of η Boo is expected to be very thin, containing less than 1% of the total mass of the star, observations

¹For preprints, please email gerard@ipac.caltech.edu.

by *MOST* were motivated by the possibility of detecting *g*-modes, along with the *p*-modes indicative of turbulent convection. In detecting eight consecutive low-frequency radial *p*-modes for this G0IV star, the *MOST* team was able to estimate many stellar parameters, including temperature, age, and mass. Additionally, new models examined by Straka et al. (2006) are able to match simultaneously the space- and ground-based pulsation data for η Boo through the inclusion of turbulence in the stellar models.

Using interferometry to obtain a *direct*, absolute measurement of this object’s linear size and effective temperature, in conjunction with the the mass estimate from the models fit to the *MOST* data, we may also infer its surface gravity, $\log g$. Values of $\log g$ are frequently utilized in stellar modeling and spectroscopy, and a direct characterization of $\log g$ for this slightly evolved object is of significant utility. We show that the combination of the high-precision photometry of *MOST* with the high-spatial resolution observations of the Palomar Testbed Interferometer (*PTI*) make for a potent combination for uniquely interpreting a star’s astrophysical parameters.

2. Interferometric Observations and Data Reduction

PTI is a three-element long-baseline interferometer, which combines two of its three 40-cm apertures to make measurements of fringe visibility, V^2 , for astronomical sources. These measurements are made at either *H*– (1.6 μm) or *K*–band (2.2 μm) with *PTI*; for this particular investigation, the K band was employed, being spectrally dispersed into 5 ‘narrow’ bands across K, centered at 2.009, 2.106, 2.203, 2.299 and 2.396 μm . For all of these observations, *PTI*’s 109.8-m N-S baseline was utilized; details on *PTI* can be found in Colavita et al. (1999).

η Boo was observed along with the unresolved calibration sources HD117176 (70 Vir), HD120136 (τ Boo), HD121107, HD121560, on 18 nights between 2000 and 2005. In addition to η Boo, a resolved check star, HD121860, was observed as a means to monitor system performance. All of the calibrators did not exceed *PTI*’s point-like calibrator angular size criterion of $\theta_{EST} < 1.0$ mas (van Belle & van Belle 2005) for absolute measurement calibration. A previous interferometric measure of η Boo’s size was made by Thévenin et al. (2005), but resolved calibration sources used in this study (due to sensitivity limitations) makes the resulting η Boo diameter estimate subject to potential systematic errors. Two of the four calibrators, 70 Vir and τ Boo, are associated with radial velocity planets (Marcy & Butler 1996; Butler et al. 1997), but no evidence has been found for V^2 variations indicative of face-on binary stars that could supplant the RV planet interpretation for these objects. The relevant observing parameters are found in Table 1, along with parameters to be derived in

§3.1.

The calibration of the η Boo V^2 data is performed by estimating the interferometer system visibility (V_{SYS}^2) using the calibration source with a model angular diameter and then normalizing the raw η Boo visibility by V_{SYS}^2 to estimate the V^2 measured by an ideal interferometer at that epoch (Mozurkewich et al. 1991; Boden et al. 1998; van Belle & van Belle 2005). Uncertainties in the system visibility and the calibrated target visibility are inferred from internal scatter among the data in an observation using standard error-propagation calculations (Colavita 1999). Calibrating our point-like calibration objects against each other produced no evidence of systematics, with all objects delivering reduced $V^2 = 1$. The observation dates, calibrated visibilities for each wavelength bin, (u, v) coordinates, and observation hour angle are presented in Table 2 for η Boo and Table 3 for HD121860. Plots of the absolute visibility data for η Boo are found in Figure 1. *PTTs* limiting night-to-night measurement error is $\sigma_{V_{\text{SYS}}^2} \approx 1.5 - 1.8\%$, the source of which is most likely a combination of effects: uncharacterized atmospheric seeing (in particular, scintillation), detector noise, and other instrumental effects. This measurement error limit is an empirically established floor from the previous study of Boden et al. (1999).




Fig. 1.— Absolute visibility data for η Boo, as discussed in §2. The line fit to the data is the visibility function corresponding to a 2.1894 ± 0.0038 mas limb-darkened angular disk

Table 1. Results from spectral energy distribution fitting, including reddening, estimated angular size, and bolometric flux at source.

	RA (J2000)	DE (J2000)	V (mag)	K (mag)	Spectral Template	A_V (mag)	θ_{EST} (mas)	
η Boo:								
HD121370	13 54 41.002	+18 23 55.30	2.68	1.485	G0IV	0.000 ± 0.011	2.320 ± 0.072	(2)
Resolved check star:								
HD121860	13 58 01.602	+07 27 48.40	7.5	2.448	M2III	0.927 ± 0.043	2.010 ± 0.109	(2)
Calibrators:								
HD117176	13 28 25.809	+13 46 43.63	5.000	3.500	G5V	0.117 ± 0.015	0.986 ± 0.020	(3)
HD120136	13 47 15.743	+17 27 24.86	4.5	3.507	F5IV	0.214 ± 0.017	0.892 ± 0.042	(4)
HD121107	13 53 12.931	+17 55 58.33	5.711	4.077	G5III	0.002 ± 0.043	0.837 ± 0.052	(1)
HD121560	13 55 49.994	+14 03 23.41	6.1	4.843	F6V	0.105 ± 0.016	0.442 ± 0.010	(1)

3. Effective Temperature and Radius Determinations

3.1. Bolometric Flux Estimates

For each of the target and calibrator stars observed in this investigation, a spectral energy distribution (SED) fit was performed. This fit was accomplished using photometry available in the literature as the input values, with template spectra appropriate for the spectral types indicated for the stars in question. The template spectra, from Pickles (1998), were adjusted to account for overall flux level and wavelength-dependent reddening, resulting in an estimate of angular size. Reddening corrections were based upon the empirical reddening determination described by Cardelli et al. (1989), which differs little from van de Hulst’s theoretical reddening curve number 15 (Johnson 1968; Dyck et al. 1996). Both narrowband and wideband photometry in the $0.3\ \mu\text{m}$ to $30\ \mu\text{m}$ were used as available, including Johnson *UBV* (see, for example, Eggen (1963, 1972); Moreno (1971)), Strömgren *ubvy* β (Piirola 1976), Geneva (Rufener 1976), 2Mass *JHK_s* (Cutri et al. 2003), and Vilnius *UPXYZS* (Zdanavicius et al. 1972); zero-magnitude flux density calibrations were based upon the values given in Cox (2000).

Starting with a reference spectral type and luminosity class as cited by SIMBAD, template spectra were fit to the photometric data. Templates in adjacent locations in spectral type and luminosity class were also tested for best fit, with the fit with best χ^2 being selected in the end for use in this study. For example, a star indicated by SIMBAD to be a G0IV would have its photometry fit to the 9 templates of spectral type F9, G0, and G1, and for luminosity classes III, IV, and V. Metallicities for these fits were assumed to be roughly solar, which is consistent with the values found for these objects in the references listed in Cayrel de Strobel et al. (1997) and Cayrel de Strobel et al. (2001).

From the best SED fit, estimates were obtained for each star for their reddening (A_V) and bolometric flux (F_{BOL}); since effective temperature was fixed for each of the Pickles (1998) library spectra, an estimate of angular size (θ_{EST}) was also obtained. The results of the fitting are given in Table 1. As an example, the SED fitting plot for η Boo is given in Figure 2.

For our calibration sources, *a priori* estimates of their sizes are necessary to account for residual resolution that may be afforded by an interferometer’s long baselines. With an expected limb darkened size of $\theta_{\text{EST}} \leq 1.00$ mas from the SED fit, calibrators have predicted V^2 ’s of $> 86\%$ for a 110-m baseline used at $\lambda = 2.2\ \mu\text{m}$ (with $V^2 > 96\%$ expected for our smallest calibrator, HD 121560). We consider this size effectively identical to its uniform disk size, since for most of our potential calibration sources, their effective temperatures are in excess of $\sim 5000\text{K}$. The difference between the uniform disk and limb darkened sizes is at

Table 2—Continued

Date	UT	$V^2(2.009\mu\text{m})$	$V^2(2.106\mu\text{m})$	$V^2(2.203\mu\text{m})$	$V^2(2.299\mu\text{m})$	$V^2(2.396\mu\text{m})$	U (m)	V (m)	HA
4/3/2005	9.19	0.562 ± 0.021	0.500 ± 0.013	0.565 ± 0.020	0.544 ± 0.014	0.489 ± 0.017	-32.86	-99.66	0.27
4/3/2005	9.50	0.592 ± 0.021	0.548 ± 0.021	0.579 ± 0.021	0.517 ± 0.022	0.501 ± 0.019	-27.74	-100.44	0.58
4/3/2005	9.79	0.576 ± 0.036	0.495 ± 0.027	0.530 ± 0.037	0.561 ± 0.039	0.486 ± 0.022	-22.63	-101.06	0.88
4/5/2005	8.30	0.560 ± 0.018	0.503 ± 0.017	0.539 ± 0.019	0.479 ± 0.018	0.457 ± 0.022	-44.44	-97.24	-0.49
4/5/2005	8.67	0.499 ± 0.012	0.560 ± 0.015	0.470 ± 0.015	0.535 ± 0.016	0.454 ± 0.021	-39.01	-98.51	-0.12
4/5/2005	8.95	0.471 ± 0.032	0.441 ± 0.030	0.539 ± 0.033	0.515 ± 0.033	0.430 ± 0.031	-34.50	-99.38	0.17
4/5/2005	9.25	0.550 ± 0.028	0.495 ± 0.025	0.534 ± 0.034	0.476 ± 0.025	0.454 ± 0.034	-29.68	-100.16	0.46
4/5/2005	9.55	0.589 ± 0.015	0.537 ± 0.013	0.506 ± 0.018	0.573 ± 0.015	0.484 ± 0.022	-24.63	-100.83	0.76
4/5/2005	9.86	0.582 ± 0.015	0.494 ± 0.020	0.527 ± 0.016	0.567 ± 0.017	0.469 ± 0.024	-19.08	-101.41	1.08

Table 3. Calibrated visibility data obtained by PTI for HD121860.

Date	UT	$V^2(2.009\mu\text{m})$	$V^2(2.106\mu\text{m})$	$V^2(2.203\mu\text{m})$	$V^2(2.299\mu\text{m})$	$V^2(2.396\mu\text{m})$	U (m)
4/9/2000	7.83	0.479 ± 0.026	0.563 ± 0.035	0.593 ± 0.017	0.620 ± 0.037	0.621 ± 0.046	-47.91
4/9/2000	8.08	0.524 ± 0.028	0.568 ± 0.034	0.592 ± 0.018	0.605 ± 0.032	0.620 ± 0.043	-44.39
5/2/2000	6.53	0.534 ± 0.018	0.575 ± 0.014	0.602 ± 0.012	0.641 ± 0.014	0.647 ± 0.015	-44.99
5/2/2000	6.80	0.522 ± 0.018	0.565 ± 0.014	0.592 ± 0.008	0.623 ± 0.013	0.630 ± 0.012	-41.11
5/4/2000	7.06	0.556 ± 0.036	0.585 ± 0.040	0.621 ± 0.030	0.643 ± 0.040	0.640 ± 0.034	-35.04
6/12/2000	3.53	0.562 ± 0.040	0.580 ± 0.030	0.613 ± 0.032	0.634 ± 0.029	0.637 ± 0.026	-49.23
6/12/2000	4.41	0.570 ± 0.049	0.597 ± 0.039	0.626 ± 0.038	0.646 ± 0.036	0.651 ± 0.032	-36.50
3/17/2001	9.81	0.507 ± 0.078	0.538 ± 0.081	0.550 ± 0.082	0.587 ± 0.072	0.594 ± 0.063	-41.43
3/19/2001	9.01	0.506 ± 0.035	0.542 ± 0.022	0.569 ± 0.028	0.595 ± 0.026	0.612 ± 0.036	-50.66
4/28/2003	6.51	0.516 ± 0.030	0.546 ± 0.023	0.577 ± 0.021	0.599 ± 0.019	0.610 ± 0.028	-49.50
4/28/2003	6.79	0.508 ± 0.035	0.529 ± 0.030	0.558 ± 0.026	0.579 ± 0.024	0.601 ± 0.032	-45.74
4/28/2003	7.11	0.521 ± 0.037	0.559 ± 0.029	0.596 ± 0.027	0.617 ± 0.024	0.633 ± 0.031	-41.16




Fig. 2.— Spectral energy distribution fitting for η Boo, as discussed in §3.1. Vertical bars on the data points represent the photometric errors, and horizontal bars correspond to the bandpass of the photometric filter being used.

the few percent level (Davis et al. 2000; Claret & Hauschildt 2003), which is far less than our size estimate error or, in particular, its impact upon the system visibility estimate derived from observations of our calibrators. A $\leq 5\%$ uncertainty in angular size will contribute, at most, less than $\leq 1.3\%$ uncertainty to the system visibility V_{sys}^2 for *PTI*. The night-to-night limiting measurement error of *PTI* is $\sigma_{V_{\text{sys}}^2} \approx 1.5 - 1.8\%$ (Boden et al. 1999), any measures of V^2 using our calibrators will be free from any potential bias in its angular size measurement at the $\sigma_\theta/\theta \approx 7\%$ level for our largest calibrator, HD117176, and at better levels of insensitivity for our the smaller calibrators.

3.2. Angular Sizes of η Boo and HD 121860

We may fit our observed visibility data to a uniform disk approximation to get an initial estimate of the angular sizes of η Boo and HD 121860. From the $V^2 = [2J_1(x)/x]^2$, where $x = \pi B\theta_{UD}\lambda^{-1}$, we get angular sizes θ_{UD} of 2.1528 ± 0.0037 mas and 2.035 ± 0.009 mas for η Boo and HD 121860, respectively, with reduced χ^2 values of 0.90 and 0.48. Given η Boo’s low rotation value of 13.5 km/s (Reiners 2006), rotational oblateness did not need to be considered in the fit (van Belle et al. 2001).

For limb darkened fits, we utilized the visibility function for a limb-darkened stellar disk as parameterized with a linear limb-darkening coefficient, μ_λ (Hanbury Brown et al. 1974):

$$V^2 = \left(\frac{1 - \mu_\lambda}{2} + \frac{\mu_\lambda}{3} \right)^{-2} \times \left[(1 - \mu_\lambda) \frac{J_1(x)}{x} + \mu_\lambda \frac{j_1(x)}{x} \right]^2 \quad (1)$$

where $x = \pi B\theta_{LD}\lambda^{-1}$. For these fits we used the $2.2 \mu\text{m}$ coefficients of $\mu_\lambda = 0.22$ and 0.38 for η Boo and HD 121860, respectively (Claret et al. 1995). Examination of the linear limb darkening coefficients from Claret et al. (1995) indicate that the value of $\mu_\lambda = 0.22 \pm 0.02$ is sufficient to account for the 5 narrowband channels of the *PTI* data (eg., μ_λ varies by less than $\Delta\mu_\lambda = 0.04$ between $2.0 \mu\text{m}$ and $2.4 \mu\text{m}$). Fitting our data, we get limb darkened angular sizes of $\theta_{LD} = 2.1894 \pm 0.0038$ mas and 2.100 ± 0.009 mas for η Boo and HD 121860, respectively, with no appreciable change in the reduced χ^2 values as compared to the uniform disk fits. A previous limb-darkened angular size of 2.25 ± 0.25 mas for η Boo was measured by Mozurkewich et al. (2003) and is consistent with our measurement.

These errors are sufficiently small that additional sources of error need to be considered. First, knowledge of *PTI*’s operational wavelength has a limit of $\sigma_\lambda \approx 0.01 \mu\text{m}$; and Second, the limb darkening coefficient μ_λ is estimated to be accurate to only $\sigma_\mu \approx 0.02$. Incorporating these effects into our LD fit for η Boo, we find an additional uncertainty contribution of 0.0040 mas, resulting in $\theta_{LD} = 2.1894 \pm 0.0055$ mas, with no appreciable increase in error for HD

121860, where the measurement error dominates the uncertainty due to the smaller number of measurements. We shall return to the estimate of uncertainty on η Boo’s angular size in §4, where the limits on our knowledge of η Boo’s possible binarity in fact ultimately limits the lower bound on our knowledge of the star’s angular size.

The absolute value of θ_{LD} is in agreement with that of the VLTI measurement in Thévenin et al. (2005), who quote $\theta_{LD} = 2.200 \pm 0.027 \pm 0.016$ mas (“statistical” and “systematic” errors are cited separately). This previous measurement is based upon limited data (only 3 V^2 data points) and is anchored to the angular size estimates of Cohen et al. (1999). Tracing back through the calibration history outlined in their paper, this is an indication that the SED angular size estimates from Cohen et al. (1999) of resolved calibrators α Crt and μ Vir they used to calibrate the system were accurate at the stated uncertainties, although this is uncertain – no values were quoted in the manuscript – and that Thévenin et al.’s measurement process preserved that SED accuracy. Additionally, no evaluation of the impact of possible binarity was considered by Thévenin et al. (2005), possibly due to the limits of their relatively small sample. Our result, since it is anchored to unresolved calibration sources, avoids the danger of being susceptible to any potential significant systematic error from SED modeling.

3.3. T_{EFF} and R

The effective temperature can be directly derived from the bolometric flux and the limb-darkening angular size:

$$T_{EFF} = 2341 \times \left[\frac{F_{BOL}}{\theta_{LD}^2} \right]^{1/4} \quad (2)$$

where F_{BOL} are in 10^{-8} ergs $\text{cm}^{-2} \text{ s}^{-1}$ and θ_{LD} is in mas (van Belle et al. 1999). Stellar radius is given by $R = 0.1076\theta_{LD}d$; where R is in $R_{\odot}$, d is in parsecs, and θ_{LD} is used as a proxy for the Rosseland mean diameter. Luminosity can be derived directly from the radius and effective temperature, $L = 4\pi R^2 \sigma T_{EFF}^4$ and is wholly independent of PTI ’s measure of θ , being depending only on d and our estimate of F_{BOL} (and, by extension, A_V). These derived values are summarized in Table 4. Our value of $L = 8.89 \pm 0.16 L_{\odot}$ is statistically consistent with the independent Guenther et al. (2005) value, indicating our F_{BOL} value discussed in §3.1 is accurate.

The measured effective temperature for HD 121860, $T_{EFF} = 3627 \pm 46$ K, which differs only slightly from the expected $T_{EFF} = 3750 \pm 22$ K (van Belle et al. 1999) for a M2III spectral type derived from the described fitting in §3.1. Its radius of $147 \pm 92 R_{\odot}$ exceeds

the expected value of $\sim 60R_{\odot}$, but the error on the radius is sufficiently large (due to the large error on the parallax) that it is not inconsistent with the smaller expected radius. The agreement of HD121860’s T_{EFF} with the expected value was an indication to us that our confidence in *PTT*s data and its error estimates are reasonable.

4. Non-Detection of Binarity of η Boo

η Boo has historically been listed as a possible spectroscopic binary (Bertiau 1957; Vansina & de Greve 1982), with an orbital confirmation being cited by Abt & Levy (1976). However, speckle interferometry observations (McAlister 1978; Bonneau et al. 1980; Hartkopf & McAlister 1984; McAlister et al. 1992) showed no evidence for detection of a possible companion to η Boo, despite an expected angular separation of $0.170''$ (Halbwachs 1981), and particularly given an quoted luminosity ratio of ≈ 0.75 (Vansina & de Greve 1982). However, the original discovery paper for η Boo’s binary nature (Bertiau 1957) suggests a late K- or M-type dwarf companion to the G0IV subgiant primary, which would indicate a brightness difference of at least $\Delta K \approx 4.5$, a substantially greater value than indicated by Vansina & de Greve (1982), and one consistent with the speckle non-detections. Abt & Levy (1976) also cite an earlier astrometric detection (Daniel & Burns 1939), although this seems unlikely given both the expected separation and the brightness ratio. The spectroscopy of Brown et al. (1997) indicates η Boo’s barycentric velocity being influenced by binarity, which Christensen-Dalsgaard et al. (1995) have suggested is indeed due to a M dwarf.

Nevertheless, given the higher resolution of *PTI*, a possible close-separation secondary companion may affect our measures of η Boo’s V^2 and thereby complicate our interpretation. As such, it was prudent for us to examine our data for evidence of ΔV^2 excursions indicative of binarity. As seen in Figure 1 and the data contained in Table 2, the η Boo V^2 data are consistent with a single star hypothesis, incorporating a wide range of (u, v) coverage and dates.

To explore to what extent a secondary star could be hidden within our data points, we examined in detail the residuals found in our single star fit, as seen in the bottom panel of Figure 1. We began by creating a synthetic V^2 dataset corresponding to a single star observed by *PTI* with $\theta_{LD} = 2.2$ mas, to which we added varying amounts of measurement noise. We then characterized the V^2 residuals through use of a histogram created through averaging two dozen runs of this synthesis, stepping through the residual values in increments of $\Delta\sigma_{V^2} = 0.01$, and comparing that histogram to that of our actual data. We found that the best fit was for measurement noise at the $\sigma_{V^2} = 0.018$ level with reduced χ^2 fit value of $\chi^2_{\nu} = 1.66$, consistent with our expectation for *PTI* data discussed in §3.1.

We then repeated this process, including measurement noise at the $\sigma_{V^2} = 0.018$ level but also adding a main sequence stellar companion of varying spectral type, using standard values of luminosity and color from Cox (2000), and constraining the orbital parameters to match the reported period of $P = 494^d$. The reduced χ^2 fit value increased as expected as the mass of the putative companion increased, but we could not exclude a possible companion of spectral type M7 and lower in a statistically meaningful way. The net result of any potential V^2 bias of a M7 companion with $\Delta K \approx 5.5$ would be to decrease the actual size of the η Boo primary by 0.014 mas. As such, we are including an additional, negative error in our calculations for the absolute parameters of η Boo to account for our uncertainty regarding this possible companion.

Given the long time baseline of V^2 measures that are present in our full dataset, it seems unlikely that chance geometric alignment would persist in making a secondary companion not appear in the form of ΔV^2 excursions. It is still entirely possible that a secondary companion is present, but at a brightness ratio that makes it not a factor in determining η Boo’s size ($L_2/L_1 < 0.005$), which is consistent with the original Bertiau (1957) result.

5. Discussion the Astrophysical Parameters of η Boo

Placing η Boo on a H-R diagram, just as was done in Guenther et al. (2005) (see their Figure 6), we may compare our values for the star’s luminosity and effective temperature to those best fit to the *MOST* data. We have highlighted this region of interest in Figure 3. We find our $\{\log L = 0.9490 \pm 0.0076, \log T = 3.7853 \pm 0.0020\}$ coordinates fall within the locus of points defined by the models the *MOST* team. The error ellipse of our $\{\log L, \log T\}$ derived from the PTI data encompasses the *MOST* $\{X = 0.71, Z = 0.04\}$ composition model point, which ultimately was favored by Guenther et al. as the best fit to their *MOST* data.

The $\{\log L, \log T\}$ coordinates defined by Di Mauro et al. (2003) were sufficiently displaced from all of the possible *MOST* coordinates that Guenther et al. suggested the Di Mauro et al. (2003) coordinates were incorrect. Our η Boo data and analysis are clearly consistent with this suggestion by Guenther et al. (2005).

5.1. Surface Gravity of η Boo

With the mass for η Boo established from the asteroseismic constraints of *MOST*, and the radius from interferometric observations, we may directly establish the surface gravity




Fig. 3.— Comparison of *MOST* fits to luminosity and effective temperature values determined in this study and by Di Mauro et al. (2003). The *MOST* fit corresponding to a model composition of $\{X = 0.71, Z = 0.04\}$ was within the error ellipse of this study’s $\{\log L = 0.9490 \pm 0.0076, \log T = 3.7853 \pm 0.0020\}$ coordinates, and was ultimately favored in Guenther et al. (2005) as the best fit to their data.

for η Boo:

$$g = \frac{GM}{R^2} \quad (3)$$

Using the *MOST* derived mass of $M = 1.71 \pm 0.05 M_{\odot}$ and the *PTI* derived radius of $R = 2.672 \pm 0.024 R_{\odot}$, we derive a surface gravity of $\log g = 3.817 \pm 0.015$ [cm s⁻²].

The *PTI* η Boo $\log g$ result is in significant disagreement with the “trigonometric” gravities found in Allende Prieto et al. (1999), who found $\log g = 3.47 \pm 0.10$ [cm s⁻²]. Malagnini & Morossi (1990) quote a value of $\log g = 3.8 \pm 0.15$ [cm s⁻²] from stellar evolution theories, although their angular size, radius and mass values on which those theories were baselined are divergent from the *PTI* and *MOST* results at 10-20% level. The study by Laird (1985) using photometric and spectrophotometric data for T_{EFF} , and Strömgren photometry plus intermediate dispersion spectra for $\log g$, is in reasonable agreement with our values, quoting $T_{EFF} = 5930 \pm 70$ K and $\log g = 3.71 \pm 0.15$ [cm s⁻²]. Similarly, Morossi et al. (2002) find $\log g = 3.71 \pm 0.13$ [cm s⁻²] using ultraviolet-visual spectrophotometry. In all of these cases, indirect measures of $\log g$ and other astrophysical parameters are overshadowed by the more direct, empirical methods afforded by the unique capabilities of *PTI* and *MOST*.

6. Conclusions

While considerably more accurate, our angular diameter determination is statistically consistent with the earlier measurement from Thévenin et al. (2005), an apparent indication that these results are free from systematic error at their stated level on uncertainty. Our angular diameter and bolometric flux values have led to an effective temperature and luminosity for the evolved star η Boo that is in direct agreement with those established by the *MOST* asteroseismology mission. Furthermore, in conjunction with the *MOST* value for stellar mass, a measure of stellar surface gravity may be made.

The combination of the precise mass estimate from *MOST* and the accurate radius measure of *PTI* has allowed us to derive a precise value of the surface gravity for η Boo. The measurement of the surface gravity η Boo, independent of spectroscopy, is a significant demonstration of the astrophysical investigative value of combining high spatial resolution interferometry with high temporal resolution photometry. Of the determinations of $\log g$ for η Boo in the literature, we find that our value has a claimed precision an order of magnitude greater than previous measures.

We gratefully acknowledge fruitful discussions with Jaymie Matthews and Theo ten Brummelaar. Science operations with PTI are conducted through the efforts of the PTI Col-

laboration (<http://pti.jpl.nasa.gov/ptimembers.html>), and we acknowledge the invaluable contributions of our PTI colleagues. We particularly thank Kevin Rykoski for his professional operation of PTI. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. Funding for PTI was provided to the Jet Propulsion Laboratory under its TOPS (Towards Other Planetary Systems), ASEPS (Astronomical Studies of Extrasolar Planetary Systems), and Origins programs and from the JPL Director's Discretionary Fund. Portions of this work were performed at the Jet Propulsion Laboratory, California Institute of Technology under contract with the National Aeronautics and Space Administration.

REFERENCES

- Abt, H. A., & Levy, S. G. 1976, *ApJS*, 30, 273
- Allende Prieto, C., García López, R. J., Lambert, D. L., & Gustafsson, B. 1999, *ApJ*, 527, 879
- Bertiau, F. C. 1957, *ApJ*, 125, 696
- Boden, A. F., Colavita, M. M., van Belle, G. T., & Shao, M. 1998, *Proc. SPIE*, 3350, 872
- Boden, A. F., et al. 1999, *ApJ*, 515, 356
- Bonneau, D., Blazit, A., Foy, R., & Labeyrie, A. 1980, *A&AS*, 42, 185
- Brown, T. M., Kennelly, E. J., Korzennik, S. G., Nisenson, P., Noyes, R. W., & Horner, S. D. 1997, *ApJ*, 475, 322
- Butler, R. P., Marcy, G. W., Williams, E., Hauser, H., & Shirts, P. 1997, *ApJ*, 474, L115
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Cayrel de Strobel, G., Soubiran, C., & Ralite, N. 2001, *A&A*, 373, 159
- Cayrel de Strobel, G., Soubiran, C., Friel, E. D., Ralite, N., & Francois, P. 1997, *A&AS*, 124, 299
- Christensen-Dalsgaard, J., Bedding, T. R., & Kjeldsen, H. 1995, *ApJ*, 443, L29
- Claret, A., Diaz-Cordoves, J., & Gimenez, A. 1995, *A&AS*, 114, 247
- Claret, A., & Hauschildt, P. H. 2003, *A&A*, 412, 241

- Cohen, M., Walker, R. G., Carter, B., Hammersley, P., Kidger, M., & Noguchi, K. 1999, *AJ*, 117, 1864
- Colavita, M. M. 1999, *PASP*, 111, 111
- Colavita, M. M., et al. 1999, *ApJ*, 510, 505
- Cox, A. N. 2000, *Allen’s astrophysical quantities*, 4th ed. Publisher: New York: AIP Press; Springer, 2000. Edited by Arthur N. Cox. ISBN: 0387987460,
- Cutri, R. M., et al. 2003, *VizieR Online Data Catalog*, 2246,
- Daniel, Z., and Burns, K., 1939, *Pub. A.A.S.*, 9, 146
- Davis, J., Tango, W. J., & Booth, A. J. 2000, *MNRAS*, 318, 387
- Di Mauro, M. P., Christensen-Dalsgaard, J., Kjeldsen, H., Bedding, T. R., & Patern, L. 2003, *A&A*, 404, 341
- Dyck, H. M., Benson, J. A., van Belle, G. T., & Ridgway, S. T. 1996, *AJ*, 111, 1705
- Eggen, O. J. 1963, *AJ*, 68, 483
- Eggen, O. J. 1972, *ApJ*, 175, 787
- Guenther, D. B., et al. 2005, *ApJ*, 635, 547
- Halbwachs, J. L. 1981, *A&AS*, 44, 47
- Hanbury Brown, R., Davis, J., Lake, R. J. W., & Thompson, R. J. 1974, *MNRAS*, 167, 475
- Hartkopf, W. I., & McAlister, H. A. 1984, *PASP*, 96, 105
- Johnson, H. L., 1968, in *Stars and Stellar Systems*, 7, *Nebulae and Interstellar Matter*, ed. B.M. Middlehurst & L.H. Aller (Chicago: Univ. of Chicago Press), chap. 5
- Kjeldsen, H., Bedding, T. R., Viskum, M., & Frandsen, S. 1995, *AJ*, 109, 1313
- Kjeldsen, H., et al. 2003, *AJ*, 126, 1483
- Laird, J. B. 1985, *ApJS*, 57, 389
- Malagnini, M. L., & Morossi, C. 1990, *A&AS*, 85, 1015
- Marcy, G. W., & Butler, R. P. 1996, *ApJ*, 464, L147

- McAlister, H. A. 1978, *PASP*, 90, 288
- McAlister, H. A., Hartkopf, W. I., & Mason, B. D. 1992, *AJ*, 104, 1961
- Moreno, H. 1971, *A&A*, 12, 442
- Morossi, C., Di Marcantonio, P., Franchini, M., Malagnini, M. L., & Chavez, M. 2002, *ApJ*, 577, 377
- Mozurkewich, D., et al. 1991, *AJ*, 101, 2207
- Mozurkewich, D., et al. 2003, *AJ*, 126, 2502
- Perryman, M. A. C., et al. 1997, *A&A*, 323, L49
- Pickles, A. J. 1998, *PASP*, 110, 863
- Pirola, V. 1976, Observatory and Astrophysics Laboratory University of Helsinki Report, 1
- Reiners, A. 2006, *A&A*, 446, 267
- Rufener, F. 1976, *A&AS*, 26, 275
- Straka, C. W., Demarque, P., Guenther, D. B., Li, L., & Robinson, F. J. 2006, *ApJ*, 636, 1078
- Thévenin, F., Kervella, P., Pichon, B., Morel, P., di Folco, E., & Lebreton, Y. 2005, *A&A*, 436, 253
- van Belle, G. T., et al. 1999, *AJ*, 117, 521
- van Belle, G. T., Ciardi, D. R., Thompson, R. R., Akeson, R. L., & Lada, E. A. 2001, *ApJ*, 559, 1155
- van Belle, G. T., & van Belle, G. 2005, *PASP*, 117, 1263
- Vansina, F., & de Greve, J. P. 1982, *Ap&SS*, 87, 377
- Walker, G., et al. 2003, *PASP*, 115, 1023
- Zdanavicius, K., et al. 1972, *Vilnius Astronomijos Observatorijos Biuletenis*, 34, 3

Table 4. Summary of stellar fundamental parameters measured for η Boo and HD121860. The error on θ_{LD} includes the uncertainty discussed in §4; subsequent columns propagate the larger of the two error bars where appropriate.

Star	θ_{LD} (mas)	F_{BOL} (10^{-8} erg cm $^{-2}$ s $^{-1}$)	π^a (mas)	T_{EFF} (K)	R ($R_{\odot}$)	L ($L_{\odot}$)
η Boo	$2.1894^{+0.0055}_{-0.0140}$	221 ± 2.82	88.17 ± 0.75	6100 ± 28	2.672 ± 0.028	8.89 ± 0.16
HD121860	2.100 ± 0.009	25.4 ± 1.26	1.54 ± 0.97	3627 ± 46	147 ± 92	3350 ± 2989

^aPerryman et al. (1997)